

2:1 BEEF CATTLE BREEDER CHELATED PREMIX WITH SALT AND MONENSIN

Formula Code: 323261M

Net weight: 25 kg

Medicated with 2,200 mg/kg (0.22%) monensin sodium as an aid in the prevention of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* in cattle.

This feed contains added selenium at 30.0 mg/kg.

A list of the ingredients used in this feed may be obtained from the manufacturer.

Product contains 25% chelated minerals.

GUARANTEED ANALYSIS:

Calcium (actual)	16.00%	Phosphorus (actual)	8.00%
Sodium (actual)	8.00%	Magnesium (actual)	3.00%
Salt (actual)	20.28%		
Copper (actual)	2,800 mg/kg	Zinc (actual)	7,000 mg/kg
Iron (actual)	3,705 mg/kg	Manganese (actual)	5,000 mg/kg
Iodine (actual)	150 mg/kg	Cobalt (act)	50 mg/kg
Vitamin A (min)	800,000 iu/kg	Vitamin E (min)	3500 iu/kg
Vitamin D (min)	80,000 iu/kg	Flourine, (min)	600 mg/kg

Mixing & Feeding Directions:

Thoroughly mix 10 kg of this 2-1 Beef Cattle Breeder Monensin Premix with 952 kg of barley, 26 kg of soybean meal, 11 kg of calcium and 1 kg of Dyna K to make one tonne of beef cattle ration. Do not feed undiluted. Feed this ration to beef cattle according to the table below to provide 22 mg/kg of monensin in the complete diet. Ensure access to clean water at all times. Provide free choice good quality forage. Contact a Nutrition Partners representative for more information.

Body Weight (kg)	Dry Mater Intake (% of Body Weight)	Mixed Feed Intake (kg/day)	Daily Intake of Monensin (mg/head/day)
545	13.6 kg (2.5%)	13.6 kg	299
568	14.2 kg (2.5%)	14.2 kg	312
593	14.8 kg (2.5%)	14.8 kg	326
613	15.3 kg (2.5%)	15.3 kg	337

Warning:

1. Do not supplement monensin from other sources (e.g. other feedstuffs containing monensin or slow release devices containing monensin).
2. Keep out of reach of children.

Caution:

1. Do not exceed recommended levels as reduced average daily gains may result.
2. Do not allow dogs, horses, or other equines or guinea fowl access to formulations containing monensin. Ingestion of monensin by these species has been fatal.
3. Do not use monensin-medicated feed for the treatment of outbreaks of coccidiosis.
4. May be used in feeds containing the following pellet-binding agents: bentonite (2%), attapulgitte (2%), kaolin (2.5%), lignin sulfonate (4%), carboxymethylcellulose (0.1%), or Agri-Colloid.
5. Directions for use must be carefully followed.
6. Do not use in association with another feed containing supplemental selenium.
7. Do not feed this mineral premix to sheep.

Individual results from the use of this product may vary due to management, environmental, genetic, health and sanitation differences. Therefore, Nutrition Partners does not warrant or guarantee individual results.

**Manufactured by:
Nutrition Partners Inc
20 East Lake Ave Airdrie, AB**

403-912-0735